

Homology operations in the topological cyclic homology of a point

HÅKON SCHAD BERGSAKER
JOHN ROGNES

We consider the commutative \mathbb{S} -algebra given by the topological cyclic homology of a point. The induced Dyer–Lashof operations in mod p homology are shown to be non-trivial for $p = 2$, and an explicit formula is given. As a part of the calculation, we are led to compare the fixed point spectrum \mathbb{S}^G of the sphere spectrum and the algebraic K -theory spectrum of finite G -sets, as structured ring spectra.

[55S12](#); [19D55](#), [55P43](#), [55P92](#)

Introduction

Let $A(\star) = K(\mathbb{S})$ denote Waldhausen’s algebraic K -theory of a point [23]. It is a commutative \mathbb{S} -algebra, in the sense of Elmendorf, Kriz, Mandell and May [7], and the algebraic K -theory $A(X)$ of any space X , or more generally the algebraic K -theory $K(R)$ of any \mathbb{S} -algebra R , is a module spectrum over it. Hence it makes sense to carefully study the commutative \mathbb{S} -algebra structure of $A(\star)$, or equivalently its structure as an E_∞ ring spectrum. To the eyes of mod p homology, the primary incarnation of this structure is the Pontryagin algebra structure on $H_*(A(\star))$, together with the multiplicative Dyer–Lashof operations $Q^i: H_*(A(\star)) \rightarrow H_{*+i}(A(\star))$, as defined by Bruner, May, McClure and Steinberger [2]. Here and elsewhere we write $H_*(E)$ for the mod p homology $H_*(E; \mathbb{F}_p)$ of a spectrum E .

The additive structure of $H_*(A(\star))$ is known for $p = 2$ and for p an odd regular prime, by the second author’s papers [18] and [19], but at present the Pontryagin product and Dyer–Lashof operations are not known for this E_∞ ring spectrum. There is, however, a very good approximation to Waldhausen’s algebraic K -theory, given by the cyclotomic trace map to the topological cyclic homology of Bökstedt, Hsiang and Madsen [1]. This is a natural map $trc: K(R) \rightarrow TC(R; p)$, which we write as $trc: A(\star) \rightarrow TC(\star; p)$

in the special case when $R = \mathbb{S}$, where $TC(\star; p) = TC(\mathbb{S}; p)$ is the topological cyclic homology of a point. By a theorem of Dundas [5], there is a homotopy cartesian square

$$\begin{array}{ccc} A(\star) & \longrightarrow & K(\mathbb{Z}) \\ \downarrow \text{trc} & & \downarrow \text{trc} \\ TC(\star; p) & \longrightarrow & TC(\mathbb{Z}; p) \end{array}$$

(after p -adic completion) of commutative \mathbb{S} -algebras [9, Sect. 6], and this square is the basis for our additive understanding of $H_*(A(\star))$.

We are therefore led to study the commutative \mathbb{S} -algebra structure of $TC(\star; p)$, including the Pontryagin algebra structure and the Dyer–Lashof operations on its mod p homology. Like in the case of algebraic K -theory, the topological cyclic homology $TC(X; p)$ of any space X , and more generally the topological cyclic homology $TC(R; p)$ of any \mathbb{S} -algebra R , is a module spectrum over $TC(\star; p)$, and this provides a second motivation for the study of $TC(\star; p)$. In the present paper, we determine the Dyer–Lashof operations in $H_*(TC(\star; p))$ in the case when $p = 2$, as explained in Theorem 0.2 and Corollary 0.3 below.

A third motivation stems from ideas of Jack Morava [17], to the effect that there may be a spectral enrichment of the algebro-geometric category of mixed Tate motives, given by A -theoretic [24] or TC -theoretic [6] correspondences, followed by stabilization. The trace map $A(\star) \rightarrow TC(\star; p) \rightarrow THH(\star) = \mathbb{S}$ defines a fiber functor to the category of \mathbb{S} -modules, with Tannakian automorphism group realized through its Hopf algebra of functions, which will be of the form $\mathbb{S} \wedge_{A(\star)} \mathbb{S}$ or $\mathbb{S} \wedge_{TC(\star; p)} \mathbb{S}$. Rationally, this is well compatible with Deligne’s results on the Tannakian group of mixed Tate motives over the integers [4]. A calculational analysis of the commutative \mathbb{S} -algebras $\mathbb{S} \wedge_{A(\star)} \mathbb{S}$ or $\mathbb{S} \wedge_{TC(\star; p)} \mathbb{S}$ clearly depends heavily on a proper understanding of the commutative \mathbb{S} -algebra structures of $A(\star)$ and $TC(\star; p)$.

Let \mathbb{T} be the circle group and let $C_{p^n} \subset \mathbb{T}$ be the (cyclic) subgroup of order p^n . The spectrum $TC(\star; p)$ is defined as the homotopy inverse limit of a diagram

$$(0-1) \quad \cdots \xrightarrow[\underset{F}{\rightrightarrows}]{R} \mathbb{S}^{C_{p^{n+1}}} \xrightarrow[\underset{F}{\rightrightarrows}]{R} \mathbb{S}^{C_{p^n}} \xrightarrow[\underset{F}{\rightrightarrows}]{R} \cdots \xrightarrow[\underset{F}{\rightrightarrows}]{R} \mathbb{S}^{C_p} \xrightarrow[\underset{F}{\rightrightarrows}]{R} \mathbb{S}$$

of E_∞ ring spectra, where $\mathbb{S}^{C_{p^n}}$ denotes the C_{p^n} -fixed points of the \mathbb{T} -equivariant sphere spectrum, the maps labeled R are restriction maps, and the maps labeled F are Frobenius maps. See Bökstedt, Hsiang and Madsen [1] or Hesselholt and Madsen [10] for the construction of these maps. Similarly, let $TC^{(1)}(\star; p)$ denote the homotopy limit of the subdiagram

$$(0-2) \quad \mathbb{S}^{C_p} \xrightarrow[\underset{F}{\rightrightarrows}]{R} \mathbb{S} ,$$

that is, the homotopy equalizer of R and F . The canonical maps

$$(0-3) \quad TC(\star; p) \xrightarrow{f_1} TC^{(1)}(\star; p) \xrightarrow{g_1} \mathbb{S}^{C_p}$$

are then maps of E_∞ ring spectra.

The unit $\eta: \mathbb{S} \rightarrow TC(\star; p)$ and the restriction $R: \mathbb{S}^{C_p} \rightarrow \mathbb{S}$ let us split off a copy of \mathbb{S} from each term in (0-3). Let $\mathbb{C}P_{-1}^\infty$ be the Thom spectrum of the negative tautological complex line bundle $-\gamma_{\mathbb{C}}^1$ over $\mathbb{C}P^\infty$. Its suspension $\Sigma\mathbb{C}P_{-1}^\infty$ is equivalent to the homotopy fiber of the dimension-shifting \mathbb{T} -transfer map $t_{\mathbb{T}}: \Sigma^\infty\Sigma(\mathbb{C}P_+^\infty) \rightarrow \mathbb{S}$, see Knapp [13, 2.9] or Lemma 1.1 below. We define the spectrum L_{-1}^∞ to be the homotopy fiber of the C_p -transfer $t_p: \Sigma^\infty(BC_p)_+ \rightarrow \mathbb{S}$. For $p = 2$, there is an equivalence $L_{-1}^\infty \simeq \mathbb{R}P_{-1}^\infty$, where $\mathbb{R}P_{-1}^\infty$ is the Thom spectrum of the negative tautological real line bundle $-\gamma_{\mathbb{R}}^1$ over $\mathbb{R}P^\infty$. The mod p homology groups of these spectra are well known:

$$\begin{aligned} H_*(\Sigma\mathbb{C}P_{-1}^\infty) &\cong \mathbb{F}_p\{\Sigma\beta_k \mid k \geq -1\} \\ H_*(L_{-1}^\infty) &\cong \mathbb{F}_p\{\alpha_k \mid k \geq -1\} \\ H_*(\Sigma^\infty(BC_p)_+) &\cong \mathbb{F}_p\{\alpha_k \mid k \geq 0\} \end{aligned}$$

Here $\Sigma\beta_k$ has degree $2k + 1$ and α_k has degree k .

Lemma 0.1 *After p -completion, diagram (0-3) is homotopy equivalent to a diagram*

$$\mathbb{S} \vee \Sigma\mathbb{C}P_{-1}^\infty \xrightarrow{1 \vee f} \mathbb{S} \vee L_{-1}^\infty \xrightarrow{1 \vee g} \mathbb{S} \vee \Sigma^\infty(BC_p)_+.$$

In particular, the Pontryagin product on $H_(TC(\star; p))$ is trivial.*

Applying homology gives a sequence

$$H_*(\mathbb{S}) \oplus H_*(\Sigma\mathbb{C}P_{-1}^\infty) \xrightarrow{1 \oplus f_*} H_*(\mathbb{S}) \oplus H_*(L_{-1}^\infty) \xrightarrow{1 \oplus g_*} H_*(\mathbb{S}) \oplus H_*(\Sigma^\infty(BC_p)_+).$$

Here f_ sends $\Sigma\beta_k$ to α_{2k+1} for $k \geq -1$, and g_* is the identity on α_k for $k \geq 0$, while α_{-1} maps to zero.*

We now state our main result, which concerns the Dyer–Lashof operations in the mod p spectrum homology $H_*(TC(\star; p))$ for $p = 2$. The calculations will be done in the auxiliary E_∞ ring spectra \mathbb{S}^{C_2} and $TC^{(1)}(\star; 2)$.

Theorem 0.2 *The Dyer–Lashof operations Q^i in $H_*(TC^{(1)}(\star; 2))$ and $H_*(\mathbb{S}^{C_2})$ are given by the formula*

$$Q^i(\alpha_j) = \binom{2^N + i - 1}{2^N + j} \alpha_{i+j},$$

where $j \geq -1$ and i is any integer, and N is sufficiently large.

Corollary 0.3 *The Dyer–Lashof operations Q^{2i} in $H_*(TC(\star; 2))$ are given by the formula*

$$Q^{2i}(\Sigma\beta_j) = \binom{2^N + i - 1}{2^N + j} \Sigma\beta_{i+j},$$

where $j \geq -1$ and i is any integer, and N is sufficiently large. The operations Q^{2i+1} are all zero for degree reasons.

Note that the binomial coefficients used in the theorem and corollary can be evaluated to

$$\binom{2^N + i - 1}{2^N + j} \equiv \begin{cases} \binom{i-1}{j} & \text{for } i > j \geq 0 \\ 1 & \text{for } (i, j) = (0, -1) \\ 0 & \text{otherwise} \end{cases}$$

modulo 2, for all sufficiently large N . In particular $Q^0(\alpha_{-1}) = \alpha_{-1}$ and $Q^0(\Sigma\beta_{-1}) = \Sigma\beta_{-1}$.

We prove Lemma 0.1 in Section 1 and Theorem 0.2 in Section 3, after a homological comparison of E_∞ ring structures in Section 2. Corollary 0.3 follows immediately from the lemma and the theorem.

1 Topological cyclic homology of a point

In this preliminary section we review the calculation of $TC(\star; p)$ from Bökstedt, Hsiang and Madsen [1, 5.17], in order to describe the map f_1 to $TC^{(1)}(\star; p)$.

For each $n \geq 1$ the Segal–tom Dieck splitting tells us that the norm–restriction homotopy cofiber sequence $\Sigma^\infty(BC_{p^n})_+ \xrightarrow{N} \mathbb{S}^{C_{p^n}} \xrightarrow{R} \mathbb{S}^{C_{p^{n-1}}}$ is canonically split. The homotopy limit $TR(\star; p) = \text{holim}_{n,R} \mathbb{S}^{C_{p^n}}$ of the R -maps in (0–1) thus factors as $TR(\star; p) \simeq \prod_{n \geq 0} \Sigma^\infty(BC_{p^n})_+$. Let $pr_n: TR(\star; p) \rightarrow \Sigma^\infty(BC_{p^n})_+$ denote the n -th projection, and let $\widetilde{TR}(\star; p) \simeq \prod_{n \geq 1} \Sigma^\infty(BC_{p^n})_+$ be the homotopy fiber of pr_0 . There is a vertical map of horizontal homotopy fiber sequences

$$(1-1) \quad \begin{array}{ccccc} TC(\star; p) & \xrightarrow{\pi} & TR(\star; p) & \xrightarrow{F-R} & TR(\star; p) \\ f_1 \downarrow & & \downarrow p_1 & & \downarrow pr_0 \\ TC^{(1)}(\star; p) & \xrightarrow{g_1} & \mathbb{S}^{C_p} & \xrightarrow{F-R} & \mathbb{S} \end{array}$$

and the augmentation $TC(\star; p) \rightarrow \mathbb{S}$ factors as $R \circ p_1 \circ \pi = pr_0 \circ \pi$. Replacing the left hand square by the homotopy fibers of the augmentations to \mathbb{S} , we get a second vertical map of horizontal homotopy fiber sequences

$$(1-2) \quad \begin{array}{ccccc} \widetilde{TC}(\star; p) & \longrightarrow & \widetilde{TR}(\star; p) & \xrightarrow{T-I} & TR(\star; p) \\ f_2 \downarrow & & \downarrow pr_1 & & \downarrow pr_0 \\ \widetilde{TC}^{(1)}(\star; p) & \xrightarrow{g_2} & \Sigma^\infty(BC_p)_+ & \xrightarrow{t_p} & \mathbb{S}. \end{array}$$

In the upper row we have used that $F - R$ restricted along the inclusion $I: \widetilde{TR}(\star; p) \rightarrow TR(\star; p)$ is homotopic to $T - I$, where T is the product of the C_p -transfer maps $\Sigma^\infty(BC_{p^n})_+ \rightarrow \Sigma^\infty(BC_{p^{n-1}})_+$ for all $n \geq 1$. See [1, (5.18)]. In the lower row we have used that $F - R$ restricted along $N: \Sigma^\infty(BC_p)_+ \rightarrow \mathbb{S}^{C_p}$ is homotopic to the C_p -transfer map t_p .

There is a third vertical map of horizontal homotopy fiber sequences

$$(1-3) \quad \begin{array}{ccccc} \text{holim}_n \Sigma^\infty(BC_{p^n})_+ & \longrightarrow & TR(\star; p) & \xrightarrow{T-1} & TR(\star; p) \\ f_3 \downarrow & & \downarrow p_1 & & \downarrow pr_0 \\ \text{hofib}(F - 2R) & \xrightarrow{g_3} & \mathbb{S}^{C_p} & \xrightarrow{F-2R} & \mathbb{S}. \end{array}$$

Replacing its left hand square by the homotopy fibers of the augmentations to \mathbb{S} , we also recover diagram (1-2).

Lemma 1.1 *There are equivalences*

$$\widetilde{TC}(\star; p) \simeq \text{hofib}(t_{\mathbb{T}}: \Sigma^\infty \Sigma(\mathbb{C}P_+^\infty) \rightarrow \mathbb{S}) \simeq \Sigma \mathbb{C}P_{-1}^\infty$$

(after p -completion) and

$$\widetilde{TC}^{(1)}(\star; p) \simeq \text{hofib}(t_p: \Sigma^\infty(BC_p)_+ \rightarrow \mathbb{S}) = L_{-1}^\infty.$$

When $p = 2$, $L_{-1}^\infty \simeq \mathbb{R}P_{-1}^\infty$.

Proof The dimension-shifting \mathbb{T} -transfer maps for the bundles $BC_{p^n} \rightarrow B\mathbb{T}$ induce an equivalence $\Sigma^\infty \Sigma(\mathbb{C}P_+^\infty) \simeq \text{holim}_n \Sigma^\infty(BC_{p^n})_+$ after p -completion [1, 5.15]. The augmentation $\text{holim}_n \Sigma^\infty(BC_{p^n})_+ \rightarrow \mathbb{S}$ then gets identified with $t_{\mathbb{T}}$, which implies the first claim.

There is a \mathbb{T} -equivariant homotopy cofiber sequence $S^0 \xrightarrow{z} S^{\mathbb{C}} \xrightarrow{t} \mathbb{T}_+ \wedge S^1$, where z is the zero-inclusion. The right hand map t is the Pontryagin–Thom collapse associated to the standard embedding $\mathbb{T} \subset \mathbb{C}$, as in Lewis, May and Steinberger [14, II.5.1]. The

dimension-shifting \mathbb{T} -transfer $t_{\mathbb{T}}: \Sigma^{\infty}\Sigma(\mathbb{C}P_{+}^{\infty}) \rightarrow \mathbb{S}$ for $E\mathbb{T} \rightarrow B\mathbb{T}$ is constructed as the balanced smash product

$$1 \wedge_{\mathbb{T}} \Sigma^{1-\mathbb{C}}(t): E\mathbb{T}_{+} \wedge_{\mathbb{T}} \Sigma^{1-\mathbb{C}}(S^{\mathbb{C}}) \rightarrow E\mathbb{T}_{+} \wedge_{\mathbb{T}} \Sigma^{1-\mathbb{C}}(\mathbb{T}_{+} \wedge S^1),$$

see [14, II.7.5]. Hence its homotopy fiber is $E\mathbb{T}_{+} \wedge_{\mathbb{T}} \Sigma^{1-\mathbb{C}}(S^0) \cong \Sigma\mathbb{C}P_{-1}^{\infty}$.

The proof that $\mathbb{R}P_{-1}^{\infty}$ is the homotopy fiber of t_p for $p = 2$ is essentially the same. \square

Proof of Lemma 0.1 Under the identifications of Lemma 1.1, the maps $f: \Sigma\mathbb{C}P_{-1}^{\infty} \rightarrow L_{-1}^{\infty}$ and $g: L_{-1}^{\infty} \rightarrow \Sigma^{\infty}(BC_p)_{+}$ correspond to the maps f_2 and g_2 in diagram (1–2), respectively.

The C_p -transfer map t_p induces multiplication by p on π_0 , and the zero map in mod p homology, so $\pi_{-1}(f)$ is surjective, f_* maps $\Sigma\beta_{-1}$ to α_{-1} , and g_* maps α_k to α_k for all $k \geq 0$. It remains to see that g_*f_* maps $\Sigma\beta_k$ to α_{2k+1} for $k \geq 0$. This is clear from diagram (1–3), since g_3f_3 agrees in positive degrees with the \mathbb{T} -transfer map $\Sigma^{\infty}\Sigma(\mathbb{C}P_{+}^{\infty}) \rightarrow \Sigma^{\infty}(BC_p)_{+}$, which has this behavior on homology. \square

2 Algebraic K -theory of finite G -sets

In this section we will compare the algebraic K -theory spectrum of finite G -sets with the G -fixed points of the sphere spectrum, as structured ring spectra. Before we state the result we recall some of the definitions involved.

The K -theory construction we use is that of Elmendorf and Mandell [8]. When the input category is a bipermutative category \mathcal{C} , their machine produces a symmetric spectrum $K(\mathcal{C})$, in the sense of Hovey, Shipley and Smith [11], with an action of the simplicial Barratt–Eccles operad. We will use the same notation for the geometrically realized symmetric spectrum in topological spaces, which has an action

$$\kappa_j: E\Sigma_j \times_{\Sigma_j} K(\mathcal{C})^{\wedge j} \rightarrow K(\mathcal{C})$$

of the operad $E\Sigma$ consisting of the contractible Σ_j -free spaces $E\Sigma_j$. As usual, $E\Sigma_j$ can be defined as the nerve $N\tilde{\Sigma}_j$ of the translation category $\tilde{\Sigma}_j$, for $j \geq 0$. The K -theory construction itself is somewhat involved, but all we need to know is that the zeroth space $K(\mathcal{C})_0$ is the nerve $N\mathcal{C}$ of \mathcal{C} , so the zeroth space of $E\Sigma_j \times_{\Sigma_j} K(\mathcal{C})^{\wedge j}$ is the nerve of $\tilde{\Sigma}_j \times_{\Sigma_j} \mathcal{C}^j$, and the action of $E\Sigma$ on $K(\mathcal{C})_0$ is given by the maps $\lambda_j: E\Sigma_j \times_{\Sigma_j} N\mathcal{C}^{\wedge j} \rightarrow N\mathcal{C}$ that are induced by the functors taking an object $(\sigma; a_1, \dots, a_j)$ in $\tilde{\Sigma}_j \times_{\Sigma_j} \mathcal{C}^j$ to the

object $a_{\sigma^{-1}(1)} \otimes \cdots \otimes a_{\sigma^{-1}(j)}$ in \mathcal{C} (see [8, Sect. 8]). Here \otimes denotes the product in the bipermutative structure on \mathcal{C} . Hence there is a commutative diagram

$$\begin{array}{ccc} E\Sigma_j \ltimes_{\Sigma_j} K(\mathcal{C})^{\wedge j} & \xrightarrow{\kappa_j} & K(\mathcal{C}) \\ \uparrow & & \uparrow \\ E\Sigma_j \ltimes_{\Sigma_j} (\Sigma^\infty NC)^{\wedge j} & \xrightarrow{\cong} \Sigma^\infty(E\Sigma_j \ltimes_{\Sigma_j} NC^{\wedge j}) \xrightarrow{\Sigma^\infty \lambda_j} & \Sigma^\infty NC \end{array}$$

for each $j \geq 0$.

Let G be a finite group, and let \mathcal{E}^G denote the category of finite G -sets and G -equivariant bijections. This is a symmetric bimonoidal category under disjoint union and cartesian product, taking (X, Y) to $X \coprod Y$ and $X \times Y$, respectively. We give $X \times Y$ the diagonal G -action. There is a functorially defined bipermutative category $\Phi\mathcal{E}^G$, and a natural equivalence $\mathcal{E}^G \rightarrow \Phi\mathcal{E}^G$ [16, VI.3.5]. It follows that there is a homotopy commutative diagram

$$(2-1) \quad \begin{array}{ccc} E\Sigma_j \ltimes_{\Sigma_j} K(\Phi\mathcal{E}^G)^{\wedge j} & \xrightarrow{\kappa_j} & K(\Phi\mathcal{E}^G) \\ \uparrow 1 \times \epsilon^{\wedge j} & & \uparrow \epsilon \\ E\Sigma_j \ltimes_{\Sigma_j} (\Sigma^\infty N\mathcal{E}^G)^{\wedge j} & \xrightarrow{\cong} \Sigma^\infty(E\Sigma_j \ltimes_{\Sigma_j} (N\mathcal{E}^G)^{\wedge j}) \xrightarrow{\Sigma^\infty \lambda_j} & \Sigma^\infty N\mathcal{E}^G \end{array}$$

for each $j \geq 0$, where

$$\lambda_j: E\Sigma_j \ltimes_{\Sigma_j} (N\mathcal{E}^G)^{\wedge j} \rightarrow N\mathcal{E}^G$$

is induced by the functor $\tilde{\Sigma}_j \ltimes_{\Sigma_j} (\mathcal{E}^G)^j \rightarrow \mathcal{E}^G$ that takes $(\sigma; X_1, \dots, X_j)$ to the cartesian product

$$X_{\sigma^{-1}(1)} \times (X_{\sigma^{-1}(2)} \times \cdots \times (X_{\sigma^{-1}(j-1)} \times X_{\sigma^{-1}(j)} \cdots)).$$

Let \mathcal{U} be a complete G -universe, and let \mathcal{L} denote the linear isometries operad with spaces $\mathcal{L}(j)$ consisting of linear isometries $\mathcal{U}^j \rightarrow \mathcal{U}$, where \mathcal{U}^j denotes the direct sum of j copies of \mathcal{U} . There is an action of G on each $\mathcal{L}(j)$ given by conjugation, and this gives \mathcal{L} the structure of an E_∞ G -operad in the sense of Lewis, May and Steinberger [14, VII.1.1]. The E_∞ ring structure on the G -equivariant sphere spectrum $\mathbb{S}_G = \Sigma_G^\infty S^0$ is given by an action

$$\zeta_j: \mathcal{L}(j) \ltimes_{\Sigma_j} \mathbb{S}_G^{\wedge j} \rightarrow \mathbb{S}_G$$

of this operad (where, for once, \ltimes denotes the twisted half-smash product in Lewis–May spectra). It is compatible with a corresponding action

$$\omega_j: \mathcal{L}(j) \ltimes_{\Sigma_j} Q_G(S^0)^{\wedge j} \rightarrow Q_G(S^0)$$

on the underlying infinite loop space $Q_G(S^0) = \Omega^\infty \mathbb{S}_G = \operatorname{colim}_{V \subset \mathcal{U}} \Omega^V S^V$, in the sense that the following diagram commutes.

$$(2-2) \quad \begin{array}{ccc} \mathcal{L}(j) \times_{\Sigma_j} (\Sigma^\infty Q_G(S^0))^{\wedge j} & \xrightarrow{\cong} & \Sigma^\infty (\mathcal{L}(j) \times_{\Sigma_j} Q_G(S^0)^{\wedge j}) \xrightarrow{\Sigma^\infty \omega_j} \Sigma^\infty Q_G(S^0) \\ \downarrow 1 \times \epsilon^{\wedge j} & & \downarrow \epsilon \\ \mathcal{L}(j) \times_{\Sigma_j} \mathbb{S}_G^{\wedge j} & \xrightarrow{\zeta_j} & \mathbb{S}_G \end{array}$$

Here ω_j sends an element in $\mathcal{L}(j) \times_{\Sigma_j} Q_G(S^0)^{\wedge j}$ represented by $(f; g_1, \dots, g_j)$, where $f: \mathcal{U}^j \rightarrow \mathcal{U}$ and $g_i: S^{V_i} \rightarrow S^{V_i}$, to the element represented by the composite of the following maps.

$$\mathcal{S}^{f(V_1 \oplus \dots \oplus V_j)} \xleftarrow[\cong]{f_*} \mathcal{S}^{V_1 \oplus \dots \oplus V_j} \xrightarrow{g_1 \wedge \dots \wedge g_j} \mathcal{S}^{V_1 \oplus \dots \oplus V_j} \xrightarrow[\cong]{f_*} \mathcal{S}^{f(V_1 \oplus \dots \oplus V_j)}$$

By taking G -fixed points we get the non-equivariant E_∞ ring spectrum $\mathbb{S}^G = (\mathbb{S}_G)^G$ with an action

$$\xi_j: \mathcal{L}^G(j) \times_{\Sigma_j} (\mathbb{S}^G)^{\wedge j} \rightarrow \mathbb{S}^G$$

of the non-equivariant E_∞ operad \mathcal{L}^G of G -equivariant isometries. The corresponding infinite loop space $\Omega^\infty(\mathbb{S}^G)$ is the space $Q_G(S^0)^G = \operatorname{colim}_{V \subset \mathcal{U}} (\Omega^V S^V)^G$, with the inherited \mathcal{L}^G -action

$$\eta_j: \mathcal{L}^G(j) \times_{\Sigma_j} (Q_G(S^0)^G)^{\wedge j} \rightarrow Q_G(S^0)^G.$$

Next we recall the definition of the Dyer–Lashof operations Q^i . Let $C_*(-)$ denote the cellular chains functor, from either CW complexes or CW spectra to chain complexes. Let E be a spectrum with an action of an E_∞ operad \mathcal{O} , and let W_* be the standard free C_p -resolution of \mathbb{F}_p with basis elements e_i in degree i . There is a chain map $W_* \rightarrow C_*(\mathcal{O}(p))$ lifting the identity on \mathbb{F}_p , unique up to homotopy, and we also denote the image of e_i under this map by e_i . Let $x \in H_q(E)$ be represented by a cycle $z \in C_q(E)$. Now consider the image of the cycle $e_i \otimes z^{\otimes p}$ under the map

$$C_*(\mathcal{O}(p)) \otimes_{\Sigma_p} C_*(E)^{\otimes p} \xrightarrow{\cong} C_*(\mathcal{O}(p) \times_{\Sigma_p} E^{\wedge p}) \xrightarrow{\xi_{p*}} C_*(E),$$

and denote its image in homology by $Q_i(x)$. Here ξ_p is the E_∞ structure map. Then for $p = 2$ define $Q^i(x) = 0$ when $i < q$, and

$$Q^i(x) = Q_{i-q}(x)$$

when $i \geq q$. For $p > 2$ define $Q^i(x) = 0$ when $2i < q$, and

$$Q^i(x) = (-1)^i \nu(q) \cdot Q_{(2i-q)(p-1)}(x)$$

when $2i \geq q$, where $\nu(q) = (-1)^{q(q-1)(p-1)/4} ((\frac{1}{2}(p-1))!)^q$. See Bruner, May, McClure and Steinberger [2, Ch. III] for more details.

The spectra \mathbb{S}^G and $K(\Phi\mathcal{E}^G)$ should be equivalent as E_∞ ring spectra, but we will only need the following weaker result.

Lemma 2.1 *There is an equivalence $\mathbb{S}^G \simeq K(\Phi\mathcal{E}^G)$ of spectra such that the induced isomorphism $H_*(\mathbb{S}^G) \cong H_*(K(\Phi\mathcal{E}^G))$ commutes with the Dyer–Lashof operations.*

Proof Our first goal is to construct a commutative diagram

$$(2-3) \quad \begin{array}{ccccc} & & E\Sigma_j \times_{\Sigma_j} (N\mathcal{E}^G)^{\wedge j} & \xrightarrow{\lambda_j} & N\mathcal{E}^G \\ & \nearrow^{1 \times \phi^{\wedge j}} & \uparrow \simeq & & \uparrow \simeq \\ E\Sigma_j \times_{\Sigma_j} (N\mathcal{E}_r^G)^{\wedge j} & \longrightarrow & (E\Sigma_j \times \mathcal{L}^G(j)) \times_{\Sigma_j} D_{\mathcal{U}}^{\wedge j} & \xrightarrow{\mu_j} & D_{\mathcal{U}} \\ & \searrow^{1 \times \psi^{\wedge j}} & \downarrow \simeq & & \downarrow \simeq \\ & & \mathcal{L}^G(j) \times_{\Sigma_j} C_{\mathcal{U}}^{\wedge j} & \xrightarrow{\nu_j} & C_{\mathcal{U}} \\ & & \downarrow & & \downarrow \psi \\ & & \mathcal{L}^G(j) \times_{\Sigma_j} (Q_G(S^0)^G)^{\wedge j} & \xrightarrow{\eta_j} & Q_G(S^0)^G \end{array}$$

We start by describing the space $C_{\mathcal{U}}$. Let V be an indexing space in \mathcal{U} . For each finite G -set X , consider the space $E_V(X)$ of X -tuples of distance-reducing embeddings of V in V , closed under the action of G . More precisely, this is the space of G -equivariant maps $\coprod_X V \rightarrow V$ such that the restriction to each summand $g: V \rightarrow V$ is an embedding that satisfies $|g(v) - g(w)| \leq |v - w|$ for all $v, w \in V$. Let $K_V(X)$ be the space of paths $[0, 1] \rightarrow E_V(X)$ such that the embeddings at the endpoint 0 are identities, and the embeddings at 1 have disjoint images. Now let

$$K_{\mathcal{U}}(X) = \operatorname{colim}_{V \subset \mathcal{U}} K_V(X).$$

These are G -equivariant versions of the spaces in the Steiner operad [22]. The group $\operatorname{Aut}^G(X)$ acts freely on $K_{\mathcal{U}}(X)$ by permuting the embeddings, and the space $C_{\mathcal{U}}$ is to be the disjoint union

$$C_{\mathcal{U}} = \coprod_{[X]} K_{\mathcal{U}}(X) / \operatorname{Aut}^G(X)$$

where X ranges over all isomorphism classes of finite G -sets.

The action of the operad \mathcal{L}^G on $C_{\mathcal{U}}$ is defined as follows. Let $f: \mathcal{U}^j \rightarrow \mathcal{U}$ be a G -linear isometry, and let $[g_i], 1 \leq i \leq j$, be elements in $C_{\mathcal{U}}$, represented by paths

of X_i -tuples of embeddings $g_i \in K_{\mathcal{U}}(X_i)$. Denote the component paths of embeddings that constitute g_i by g_{i,x_i} , where $x_i \in X_i$. The resulting element $\nu_j(f; [g_1], \dots, [g_j])$ in $C_{\mathcal{U}}$ is represented by an element in $K_{\mathcal{U}}(X_1 \times \dots \times X_j)$, which on the summand indexed by (x_1, \dots, x_j) is given by $f \circ (g_{1,x_1} \times \dots \times g_{j,x_j}) \circ f^{-1}$.

There is a map $C_{\mathcal{U}} \rightarrow Q_G(S^0)^G$, given by evaluating a Steiner path in $E_V(X)$ at 1 to get a G -equivariant embedding $e: \coprod_X V \rightarrow V$, and then applying a folded Pontryagin–Thom construction to obtain a G -equivariant map $q: S^V \rightarrow S^V$, which is a point in $Q_G(S^0)^G$. Given the distance-reducing embedding e , let $S^V \rightarrow \bigvee_X S^V$ be the G -equivariant map that is given by e^{-1} on the image of e in $V \subset S^V$ and maps the remainder of S^V to the base point of $\bigvee_X S^V \supset \coprod_X V$. Let $\bigvee_X S^V \rightarrow S^V$ be the fold map that is the identity on each summand. The folded Pontryagin–Thom construction q is the composite of these two G -maps. If we permute the embeddings indexed by X we get the same element in $Q_G(S^0)^G$, so our map is well-defined. A comparison of definitions shows that this construction is compatible with the \mathcal{L}^G -actions on $C_{\mathcal{U}}$ and $Q_G(S^0)^G$, so the lower square in (2–3) commutes.

Let

$$D_{\mathcal{U}} = \coprod_{[X]} (E \operatorname{Aut}^G(X) \times K_{\mathcal{U}}(X)) / \operatorname{Aut}^G(X),$$

where $\operatorname{Aut}^G(X)$ acts diagonally on the product. The nerve $N\mathcal{E}^G$ splits as a sum of components

$$(2-4) \quad N\mathcal{E}^G \simeq \coprod_{[X]} B \operatorname{Aut}^G(X),$$

where the disjoint union is over the isomorphism classes of G -sets X . Projection on the first factor in $D_{\mathcal{U}}$ followed by this homotopy equivalence gives the map $D_{\mathcal{U}} \rightarrow N\mathcal{E}^G$ in (2–3), while the map $D_{\mathcal{U}} \rightarrow C_{\mathcal{U}}$ is the projection on the second factor. There is an induced action of the product operad $E\Sigma \times \mathcal{L}^G$ on $D_{\mathcal{U}}$, defined as follows. Let $(e, f) \in E\Sigma_j \times \mathcal{L}^G(j)$ and let $(e_i, f_i) \in E \operatorname{Aut}^G(X) \times C_{\mathcal{U}}(X)$ represent elements in $D_{\mathcal{U}}$, for $1 \leq i \leq j$. The image under μ_j is the element represented by $(\lambda_j(e; e_1, \dots, e_j), \nu_j(f; f_1, \dots, f_j))$. This makes the upper and middle squares in (2–3) commute.

Let

$$N\mathcal{E}_{tr}^G = \left(\coprod_{(H)} (E \operatorname{Aut}^G(G/H) \times K_{\mathcal{U}}(G/H)) / \operatorname{Aut}^G(G/H) \right)_+,$$

where the coproduct is taken over the conjugacy classes of subgroups H of G . The map $N\mathcal{E}_{tr}^G \rightarrow D_{\mathcal{U}}$ in (2–3) is the inclusion of the components indexed by the isomorphism classes of transitive G -sets.

The maps ϕ and ψ are defined by commutativity of the right hand triangles in the diagram. We claim that the adjoints

$$(2-5) \quad \Sigma^\infty N\mathcal{E}_{tr}^G \rightarrow K(\mathcal{E}^G) \simeq K(\Phi\mathcal{E}^G)$$

$$(2-6) \quad \Sigma^\infty N\mathcal{E}_{tr}^G \rightarrow \mathbb{S}^G$$

of the maps ϕ and ψ , respectively, are both equivalences. Here $K(\mathcal{E}^G)$ is the additive K -theory spectrum of \mathcal{E}^G , with zeroth space $K(\mathcal{E}^G)_0 = N\mathcal{E}^G$, which only depends on the additive symmetric monoidal structure of \mathcal{E}^G . There is an equivalence

$$\Sigma^\infty N\mathcal{E}_{tr}^G \simeq \bigvee_{(H)} \Sigma^\infty BW_G H_+,$$

where $W_G H = N_G H / H \cong \text{Aut}^G(G/H)$ is the Weyl group of H and the wedge sum is over the conjugacy classes of subgroups of G . By Waldhausen's additivity theorem [23, 1.3.2] applied to a suitable filtration of \mathcal{E}^G according to stabilizer types, there is a splitting

$$K(\mathcal{E}^G) \simeq \bigvee_{(H)} K(\mathcal{E}(W_G H)),$$

where $\mathcal{E}(W_G H)$ is the category of finite free $W_G H$ -sets and equivariant bijections. The map (2-5) is equivalent under these identifications to the wedge sum of the maps

$$(2-7) \quad \Sigma^\infty BW_G H_+ \rightarrow K(\mathcal{E}(W_G H))$$

that are left adjoint to the inclusions $BW_G H_+ \rightarrow N\mathcal{E}(W_G H) = K(\mathcal{E}(W_G H))_0$.

The Barratt–Priddy–Quillen–Segal theorem [20, 3.6] says that each of the maps (2-7) is an equivalence, hence (2-5) is an equivalence. The map (2-6) is an equivalence by the Segal–tom Dieck splitting [14, V.11.2]. The composition of these two equivalences is the equivalence $\mathbb{S}^G \simeq K(\Phi\mathcal{E}^G)$ referred to in the statement of the lemma.

We apply the suspension spectrum functor Σ^∞ to the diagram (2-3), combine it with diagram (2-1) and the G -fixed part of (2-2), take homology, and end up with the following commutative diagram.

(2-8)

$$\begin{array}{ccccc}
 & & H_*(\Sigma_j; H_*(K(\Phi\mathcal{E}^G))^{\otimes j}) & \xrightarrow{\kappa_{j*}} & H_*(K(\Phi\mathcal{E}^G)) \\
 & \cong \nearrow & \uparrow & & \uparrow \epsilon_1 \\
 H_*(\Sigma_j; H_*(N\mathcal{E}_{tr}^G)^{\otimes j}) & \longrightarrow & H_*(\Sigma_j; \tilde{H}_*(D\mathcal{U})^{\otimes j}) & \xrightarrow{\mu_{j*}} & \tilde{H}_*(D\mathcal{U}) \\
 & \cong \searrow & \downarrow & & \downarrow \epsilon_2 \\
 & & H_*(\Sigma_j; H_*(\mathbb{S}^G)^{\otimes j}) & \xrightarrow{\xi_{j*}} & H_*(\mathbb{S}^G)
 \end{array}$$

$\begin{array}{c} \swarrow \phi_* \\ \cong \\ \searrow \psi_* \end{array}$

We need the fact that ϵ_1 and ϵ_2 have the same kernel. In fact, all summands in $\tilde{H}_*(D\mathcal{U})$ indexed by G -sets with more than one orbit map to zero under both ϵ_1 and ϵ_2 . This follows from the fact that Pontryagin products and additive Dyer–Lashof operations vanish after stabilization. More precisely, a decomposition of a G -set $X = \coprod_{i=1}^k n_i(G/H_i)$, where the H_i lie in distinct conjugacy classes, induces a factorization

$$B \operatorname{Aut}^G(X) \cong \prod_{i=1}^k B(\Sigma_{n_i} \wr W_G H_i).$$

The homology group $H_*(B(\Sigma_{n_i} \wr W_G H_i)) \subset H_*(N\mathcal{E}^G)$ is generated by $H_*(B W_G H_i)$ under iterated Pontryagin products and Dyer–Lashof operations, see Cohen, Lada and May [3, I.4.1], which all map to zero under ϵ_1 and ϵ_2 unless $k = 1$ and $n_1 = 1$.

Let $x \in H_*(K(\Phi\mathcal{E}^G))$, and let $y \in H_*(\mathbb{S}^G)$ be the element corresponding to x under $\psi_* \circ \phi_*^{-1}$, via an element $z \in H_*(N\mathcal{E}_{tr}^G)$. We need to show that the image $Q_i(x)$ of $e_i \otimes x^{\otimes p}$ under the top map corresponds, via the isomorphism, to the image $Q_i(y)$ of $e_i \otimes y^{\otimes p}$ under the bottom map. The element $e_i \otimes z^{\otimes p} \in H_*(\Sigma_p; H_*(N\mathcal{E}_{tr}^G)^{\otimes p})$ maps to an element $Q_i(z) \in \tilde{H}_*(D\mathcal{U})$, which further maps to $Q_i(x)$ and $Q_i(y)$ under ϵ_1 and ϵ_2 , respectively. Let $w \in H_*(N\mathcal{E}_{tr}^G)$ map to $Q_i(x)$ under ϕ_* . Since the maps ϵ_1 and ϵ_2 have the same kernel, the elements $Q_i(z)$ and w have the same image in $H_*(\mathbb{S}^G)$, which implies the result. \square

Remark 2.2 The additive equivalence $\mathbb{S}^G \simeq K(\mathcal{E}^G) \simeq K(\Phi\mathcal{E}^G)$ of spectra can be realized as the G -fixed part of a G -equivalence $\mathbb{S}_G \simeq K_G(\mathcal{E})$ of G -spectra, for example using Shimakawa’s construction [21] of G -equivariant K -theory spectra. Presumably this is a G -equivalence of E_∞ ring G -spectra.

3 Proof of the main theorem

Recall the E_∞ structure maps $\lambda_j: E\Sigma_j \times_{\Sigma_j} (N\mathcal{E}^G)^{\wedge j} \rightarrow N\mathcal{E}^G$. We have inclusions $B \operatorname{Aut}^G(G) \rightarrow N\mathcal{E}^G$ and $\delta: B \operatorname{Aut}^G(G^j) \rightarrow N\mathcal{E}^G$, corresponding to the summands indexed by $X = G$ and $X = G^j = G \times \cdots \times G$, respectively, in the decomposition (2–4) of $N\mathcal{E}^G$. Restricting λ_j to these summands, we have a commutative diagram

$$(3-1) \quad \begin{array}{ccc} E\Sigma_j \times_{\Sigma_j} (N\mathcal{E}^G)^{\wedge j} & \xrightarrow{\lambda_j} & N\mathcal{E}^G \\ \uparrow & & \uparrow \delta \\ E\Sigma_j \times_{\Sigma_j} B \operatorname{Aut}^G(G^j) & \xrightarrow{\cong} B(\Sigma_j \times \operatorname{Aut}^G(G)^j) \xrightarrow{B\phi} & B \operatorname{Aut}^G(G^j), \end{array}$$

where the homomorphism ϕ sends an element $(\sigma; f_1, \dots, f_j)$ in $\Sigma_j \times \text{Aut}^G(G)^j$ to the G -automorphism $f_{\sigma^{-1}(1)} \times \dots \times f_{\sigma^{-1}(j)}$ of G^j .

We write $\Sigma_j \wr \text{Aut}^G(G) \cong \Sigma_j \wr G$ for the wreath product $\Sigma_j \times \text{Aut}^G(G)^j$. The free G -set G^j splits into $k = |G|^{j-1}$ orbits, and we fix a G -equivariant bijection $G^j \cong \coprod_k G$. This induces an isomorphism $\text{Aut}^G(G^j) \cong \text{Aut}^G(\coprod_k G)$, and we also have $\text{Aut}^G(\coprod_k G) \cong \Sigma_k \wr G$. Thus we get a commutative diagram

$$(3-2) \quad \begin{array}{ccc} B(\Sigma_j \wr \text{Aut}^G(G)^j) & \xrightarrow{B\phi} & B \text{Aut}^G(G^j) \\ \cong \uparrow & & \cong \uparrow \\ B(\Sigma_j \wr G) & \xrightarrow{B\phi} & B(\Sigma_k \wr G) \end{array}$$

where we also write ϕ for the induced homomorphism $\Sigma_j \wr G \rightarrow \Sigma_k \wr G$.

Now we specialize to the case $p = 2$. First we study the Dyer–Lashof operation $Q^2: H_1(\mathbb{S}^{C_2}) \rightarrow H_3(\mathbb{S}^{C_2})$.

Lemma 3.1 *The operation Q^2 in $H_*(\mathbb{S}^{C_2})$ satisfies $Q^2(\alpha_1) = \alpha_3$.*

Proof Let $C = C_2$. By Lemma 2.1, we may instead compute Q^2 in $H_*(K(\Phi \mathcal{E}^C))$. We let $j = 2$, combine diagrams (2–1), (3–1) and (3–2), apply homology, and end up with the upper half of the diagram

$$(3-3) \quad \begin{array}{ccc} H_*(E\Sigma_2 \times_{\Sigma_2} K(\Phi \mathcal{E}^C)^{\wedge 2}) & \xrightarrow{\kappa_{2*}} & H_*(K(\Phi \mathcal{E}^C)) \\ \uparrow & & \uparrow \epsilon_* \circ \delta_* \\ H_*(B(\Sigma_2 \wr C)) & \xrightarrow{B\phi_*} & H_*(B(\Sigma_2 \wr C)) \\ \uparrow Bd_* & & \uparrow B\iota_* \\ H_*(B(\Sigma_2 \times C)) & \xrightarrow{B\psi_*} & H_*(B(C \times C)). \end{array}$$

The vertical homomorphisms in the lower square are induced by the homomorphism $d = 1 \times \Delta$ that sends (σ, x) to $(\sigma; x, x)$, and the inclusion ι of the subgroup $C \times C = C^2$ in $\Sigma_2 \wr C = \Sigma_2 \times C^2$. The homomorphism ψ is the restriction of ϕ to $\Sigma_2 \times C$. It is easily checked that ψ takes values in the subgroup $C \times C$ (since $p = 2$) and is given by $\psi(\sigma, x) = (x, \sigma x)$, using the description of ϕ given after diagram (3–1). We have $B\psi_*(e_i \otimes 1) = 1 \otimes \alpha_i$ and

$$B\psi_*(1 \otimes \alpha_j) = \Delta_*(\alpha_j) = \sum_{s+t=j} \alpha_s \otimes \alpha_t,$$

which combine to give

$$B\psi_*(e_i \otimes \alpha_j) = \sum_{s+t=j} \alpha_s \otimes (\alpha_i * \alpha_t),$$

where $*$ denotes the Pontryagin product in $H_*(BC)$ induced by the topological group multiplication $BC \times BC \rightarrow BC$. We recall that $\alpha_i * \alpha_t = \binom{i+t}{i} \alpha_{i+t}$.

By May's paper [15, 9.1] the map Bd_* is given by

$$Bd_*(e_i \otimes \alpha_j) = \sum_k e_{i+2k-j} \otimes Sq_*^k(\alpha_j) \otimes Sq_*^k(\alpha_j).$$

Recall that $Sq_*^k(\alpha_j) = \binom{j-k}{k} \alpha_{j-k}$, where Sq_*^k denotes the dual of the Steenrod operation Sq^k . In particular $Bd_*(e_1 \otimes \alpha_2) = e_1 \otimes \alpha_1 \otimes \alpha_1$, which further maps to $Q_1(\alpha_1)$ in the upper right hand corner of (3–3). But now $Q_1(\alpha_1)$ is also the image of $e_1 \otimes \alpha_2$ under $\epsilon_* \circ \delta_* \circ B\iota_* \circ B\psi_*$. Using the description of $B\psi_*$ above, we see that $Q_1(\alpha_1)$ equals

$$(3-4) \quad \sum_{s+t=2} \epsilon_* \delta_*(1 \otimes \alpha_s \otimes (\alpha_1 * \alpha_t)).$$

The map ϵ_* vanishes on decomposables with respect to the product in $H_*(N\mathcal{E}^C)$ induced by the additive symmetric monoidal structure on \mathcal{E}^C . The element $\delta_*(1 \otimes \alpha_s \otimes (\alpha_1 * \alpha_t))$ is the image of

$$\alpha_s \otimes (\alpha_1 * \alpha_t) \in H_*(B \text{Aut}^C(C) \times B \text{Aut}^C(C))$$

in $H_*(B \text{Aut}^C(C \amalg C)) \subset H_*(N\mathcal{E}^C)$ under the map induced by disjoint union, thus the only non-zero term in (3–4) is the one with $s = 0$ and $t = 2$, and $Q^2(\alpha_1) = Q_1(\alpha_1) = \epsilon_*(1 \otimes \alpha_0 \otimes (\alpha_1 * \alpha_2)) = \alpha_3$. \square

Proof of Theorem 0.2 We now turn to the operations in $H_*(TC^{(1)}(\star; 2)) = \mathbb{F}_2 \oplus H_*(\mathbb{R}P_{-1}^\infty)$. The general formula for the Q^i will follow from Lemma 3.1 and the Nishida relations, which say in particular (see [2, III.1.1]) that

$$(3-5) \quad Sq_*^{i+j+1} Q^i(\alpha_j) = \sum_k \binom{2^N - j - 1}{2^N - i - 2j - 2 + 2k} Q^{k-j-1} Sq_*^k(\alpha_j),$$

where N is sufficiently large. When $k \geq j + 2$ the element $Sq_*^k(\alpha_j)$ is zero for degree reasons, and when $k \leq j$ the fact that Q^{k-j-1} vanishes on classes in degree higher than $k - j - 1$ implies that $Q^{k-j-1} Sq_*^k(\alpha_j) = 0$. Hence the sum in (3–5) simplifies to the single term

$$Sq_*^{i+j+1} Q^i(\alpha_j) = \binom{2^N - j - 1}{2^N - i} Q^0 Sq_*^{j+1}(\alpha_j)$$

for $k = j + 1$, where N is large.

Bob Bruner has observed that

$$\binom{2^N - j - 1}{2^N - i} \equiv \binom{2^N + i - 1}{2^N + j} \pmod{2},$$

for large N . Here is a quick proof. Let x_k denote the k 'th bit in the binary expansion of a natural number x . Then $\binom{2^N - j - 1}{2^N - i} \equiv 1$ if and only if $(2^N - i)_k = 1$ implies $(2^N - j - 1)_k = 1$ for all k , and $\binom{2^N + i - 1}{2^N + j} \equiv 1$ if and only if $(2^N + j)_k = 1$ implies $(2^N + i - 1)_k = 1$ for all k . But for N large compared to i, j and k the bit $(2^N - i)_k$ is complementary to $(2^N + i - 1)_k$, and $(2^N - j - 1)_k$ is complementary to $(2^N + j)_k$, so $\binom{2^N - j - 1}{2^N - i} \equiv 1$ if and only if $\binom{2^N + i - 1}{2^N + j} \equiv 1$.

The operations Sq_*^k in $H_*(\mathbb{R}P_{-1}^\infty)$ are given by the formula

$$Sq_*^k(\alpha_j) = \binom{j - k}{k} \alpha_{j - k}.$$

This follows by the corresponding formula for $\mathbb{R}P^\infty$ and James periodicity. More precisely, a theorem of James [12] says that given $m \leq n$, there is a positive integer M , depending only on $n - m$, such that $\mathbb{R}P_{m+\ell}^{n+\ell} \simeq \Sigma^\ell \mathbb{R}P_m^n$ when ℓ is a positive multiple of 2^M . The space $\mathbb{R}P_m^n$ is the stunted projective space $\mathbb{R}P^n / \mathbb{R}P^{m-1}$. If we now define the spectrum $\mathbb{R}P_{-1}^n$ to be $\Sigma^{-\ell} \Sigma^\infty \mathbb{R}P_{\ell-1}^{n+\ell}$ for such ℓ (depending on n), we have that $\mathbb{R}P_{-1}^\infty = \text{colim}_n \mathbb{R}P_{-1}^n$. The Steenrod operations in $H_*(\mathbb{R}P_{-1}^\infty)$ can now be calculated from the operations in $H_*(\mathbb{R}P_{\ell-1}^{n+\ell})$, and the stated formula follows by noting that the relevant binomial coefficients are 2^M -periodic in the numerator.

In particular $Sq_*^{j+1}(\alpha_j) = \alpha_{-1}$ for all $j \geq -1$, and we have

$$Sq_*^{i+j+1} Q^i(\alpha_j) = \binom{2^N + i - 1}{2^N + j} Q^0(\alpha_{-1}).$$

If $Q^0(\alpha_{-1})$ were zero, it would follow that $Q^i(\alpha_j) = 0$ for all i and j , since Sq_*^{i+j+1} is an isomorphism to dimension -1 . But this contradicts Lemma 3.1. Hence $Q^0(\alpha_{-1}) = \alpha_{-1}$, and the formula stated in the theorem follows. \square

References

- [1] M Bökstedt, W-C Hsiang, and I Madsen, *The cyclotomic trace and algebraic K-theory of spaces*, Invent. Math. **111** (1993), no. 3, 465–539.
- [2] R R Bruner, J P May, J E McClure, and M Steinberger, *H_∞ ring spectra and their applications*, Lecture Notes in Mathematics, vol. 1176, Springer-Verlag, Berlin, 1986.

- [3] F R Cohen, T J Lada, and J P May, *The homology of iterated loop spaces*, Lecture Notes in Mathematics, Vol. 533, Springer-Verlag, Berlin, 1976.
- [4] P Deligne, *Le groupe fondamental de la droite projective moins trois points*, Galois groups over \mathbf{Q} (Berkeley, CA, 1987), Math. Sci. Res. Inst. Publ., vol. 16, Springer, New York, 1989, pp. 79–297.
- [5] B I Dundas, *Relative K-theory and topological cyclic homology*, Acta Math. **179** (1997), no. 2, 223–242.
- [6] B I Dundas and P A Østvær, *A bivariant Chern character*, in progress.
- [7] A D Elmendorf, I Kriz, M A Mandell, and J P May, *Rings, modules, and algebras in stable homotopy theory*, Mathematical Surveys and Monographs, vol. 47, American Mathematical Society, Providence, RI, 1997, With an appendix by M Cole.
- [8] A D Elmendorf and M A Mandell, *Rings, modules, and algebras in infinite loop space theory*, Adv. Math. **205** (2006), no. 1, 163–228.
- [9] T Geisser and L Hesselholt, *Topological cyclic homology of schemes*, Algebraic K-theory (Seattle, WA, 1997), Proc. Sympos. Pure Math., vol. 67, Amer. Math. Soc., Providence, RI, 1999, pp. 41–87.
- [10] L Hesselholt and I Madsen, *On the K-theory of finite algebras over Witt vectors of perfect fields*, Topology **36** (1997), no. 1, 29–101.
- [11] M Hovey, B Shipley, and J Smith, *Symmetric spectra*, J. Amer. Math. Soc. **13** (2000), no. 1, 149–208.
- [12] I M James, *Spaces associated with Stiefel manifolds*, Proc. London Math. Soc. **9** (1959), 115–140.
- [13] K-H Knapp, *Some applications of K-theory to framed bordism: e-invariant and transfer*, Habilitationsschrift, Bonn, 1979, <http://elpub.bib.uni-wuppertal.de/edocs/dokumente/fbc/mathematik/publikationen/knapp/habi1979>.
- [14] L G Lewis, Jr, J P May, and M Steinberger, *Equivariant stable homotopy theory*, Lecture Notes in Mathematics, vol. 1213, Springer-Verlag, Berlin, 1986, With contributions by J E McClure.
- [15] J P May, *A general algebraic approach to Steenrod operations*, The Steenrod Algebra and its Applications (Proc. Conf. to Celebrate N E Steenrod’s Sixtieth Birthday, Battelle Memorial Inst., Columbus, Ohio, 1970), Lecture Notes in Mathematics, Vol. 168, Springer, Berlin, 1970, pp. 153–231.
- [16] ———, *E_∞ ring spaces and E_∞ ring spectra*, Lecture Notes in Mathematics, Vol. 577, Springer-Verlag, Berlin, 1977, With contributions by F Quinn, N Ray, and J Tornehave.
- [17] J Morava, *A theory of base motives*, (2009), 11 pages, arXiv:0908.3124 preprint.
- [18] J Rognes, *Two-primary algebraic K-theory of pointed spaces*, Topology **41** (2002), no. 5, 873–926.

- [19] ———, *The smooth Whitehead spectrum of a point at odd regular primes*, *Geom. Topol.* **7** (2003), 155–184 (electronic).
- [20] G Segal, *Categories and cohomology theories*, *Topology* **13** (1974), 293–312.
- [21] K Shimakawa, *Infinite loop G -spaces associated to monoidal G -graded categories*, *Publ. Res. Inst. Math. Sci.* **25** (1989), no. 2, 239–262.
- [22] R Steiner, *A canonical operad pair*, *Math. Proc. Cambridge Philos. Soc.* **86** (1979), no. 3, 443–449.
- [23] F Waldhausen, *Algebraic K -theory of topological spaces. I*, *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 1, Proc. Sympos. Pure Math., XXXII, Amer. Math. Soc., Providence, R.I., 1978*, pp. 35–60.
- [24] B Williams, *Bivariant Riemann Roch theorems*, *Geometry and topology: Aarhus (1998), Contemp. Math., vol. 258, Amer. Math. Soc., Providence, RI, 2000*, pp. 377–393.

Department of Mathematics, University of Oslo, Oslo, Norway

Department of Mathematics, University of Oslo, Oslo, Norway

hakonsb@math.uio.no, rognes@math.uio.no